

# Decomposition rank of UHF-absorbing C\*-algebras

Hiroki Matui

Graduate School of Science  
Chiba University

Inage-ku, Chiba 263-8522, Japan

Yasuhiko Sato

Graduate School of Science  
Kyoto University

Sakyo-ku, Kyoto 606-8502, Japan

## Abstract

Let  $A$  be a unital separable simple C\*-algebra with a unique tracial state. We prove that if  $A$  is nuclear and quasidiagonal, then  $A$  tensored with the universal UHF-algebra has decomposition rank at most one. Then it is proved that  $A$  is nuclear, quasidiagonal and has strict comparison if and only if  $A$  has finite decomposition rank. For such  $A$ , we also give a direct proof that  $A$  tensored with a UHF-algebra has tracial rank zero.

## 1 Introduction

E. Kirchberg and W. Winter [11] introduced the notion of decomposition rank for nuclear C\*-algebras as a generalization of topological dimension. Among others, they showed that finite decomposition rank implies quasidiagonality. The concept of decomposition rank is particularly relevant for Elliott's program to classify nuclear C\*-algebras by K-theoretic data. Indeed, all nuclear stably finite C\*-algebras classified so far by their Elliott invariants have in fact finite decomposition rank. Also, this property is expected to be equivalent to other regularity properties. W. Winter [27] proved that any unital separable simple infinite-dimensional C\*-algebra  $A$  with finite decomposition rank is  $\mathcal{Z}$ -absorbing, i.e.  $A$  absorbs the Jiang-Su algebra  $\mathcal{Z}$  tensorially. M. Rørdam [22] showed that  $\mathcal{Z}$ -absorption implies strict comparison for unital simple exact C\*-algebras. In the present paper we provide the converse direction of these results under the assumption that the algebra has a unique tracial state.

**Theorem 1.1.** *Let  $A$  be a unital separable simple C\*-algebra with a unique tracial state. If  $A$  is nuclear, quasidiagonal and has strict comparison, then the decomposition rank of  $A$  is at most three.*

The notion of strict comparison was first introduced by B. Blackadar [2] for projections, which was called FCQ2. Later this definition is extended to positive elements in the following sense. Throughout this paper, for a C\*-algebra  $A$  we denote by  $T(A)$  the set of tracial states on  $A$ . We say that  $A$  has *strict comparison* if for two positive elements  $a, b$  in  $M_k(A)$  with  $\lim_{n \rightarrow \infty} \tau(a^{1/n}) < \lim_{n \rightarrow \infty} \tau(b^{1/n})$  for any  $\tau \in T(A)$  there exist  $r_n \in M_k(A)$ ,  $n \in \mathbb{N}$  such that  $r_n^* b r_n \rightarrow a$ , where  $\tau$  is regarded as an unnormalized trace on  $M_k(A)$ . The precise definition of decomposition rank is given in Definition 4.1 and the proof of the main theorem is given in Section 5.

In 2008, A. S. Toms and W. Winter conjectured that properties of finite decomposition rank,  $\mathcal{Z}$ -absorption, and strict comparison are equivalent for unital separable simple infinite-dimensional finite nuclear  $C^*$ -algebras (see [27, Conjecture 0.1] for example). The theorem above, together with the results in [11, 22, 27, 17], gives a partial affirmative answer to this conjecture in the following sense.

**Corollary 1.2.** *Let  $A$  be a unital separable simple infinite-dimensional nuclear  $C^*$ -algebra with a unique tracial state. Then the following are equivalent.*

- (i)  *$A$  has finite decomposition rank.*
- (ii)  *$A$  is quasidiagonal and is  $\mathcal{Z}$ -absorbing.*
- (iii)  *$A$  is quasidiagonal and has strict comparison.*

The technique that we develop in proving Theorem 1.1 also allows us to bound other ranks of nuclear  $C^*$ -algebras. In Section 6, for  $A$  as in Theorem 1.1, we show that  $A$  tensored with a UHF-algebra has tracial rank zero. This enables us to remove the technical condition from H. Lin's result [13, Corollary 5.10] for nuclear  $C^*$ -algebras, and to provide a partial answer to his conjecture mentioned in the introduction of [13]. In Section 7, we show that any Kirchberg algebra has nuclear dimension at most three. We remark that W. Winter and J. Zacharias [28, Theorem 7.5] proved that if  $A$  is a Kirchberg algebra satisfying the Universal Coefficient Theorem (UCT), then it has finite nuclear dimension.

## 2 Murray-von Neumann equivalence

A celebrated result of A. Connes says that injective factors with separable predual are approximately finite dimensional (AFD), see [6]. The proof relies on his deep study of automorphisms of factors [4, 5]. By now, two alternative proofs which do not use automorphisms are known. One is due to U. Haagerup [7]. His proof uses trace preserving completely positive maps instead of automorphisms. S. Popa gave another short proof by using excision of amenable traces [20]. In the present paper, we focus on the proofs by Connes and Haagerup, in which the Murray-von Neumann equivalence for projections plays an essential role. They constructed approximation by finite dimensional subalgebras by using partial isometries inducing the Murray-von Neumann equivalence for certain projections. In this section we establish a generalization of the Murray-von Neumann equivalence (Lemma 2.1). In our setting, the 'equivalence' is given by two contractions. These two contractions will become involved in the estimate of decomposition rank of UHF-absorbing  $C^*$ -algebras in Section 4.

We say that a  $C^*$ -algebra  $A$  has *strict comparison for projections* if for any two projections  $p$  and  $q$  in  $M_k(A)$  satisfying  $\tau(p) < \tau(q)$  for any tracial state  $\tau$  of  $A$ , there exists a partial isometry  $v$  in  $M_k(A)$  such that  $v^*v = p$  and  $vv^* \leq q$ .

**Lemma 2.1.** *Let  $A$  be a unital  $C^*$ -algebra with strict comparison for projections. Let  $p, q$  be projections in  $A$  and  $n \in \mathbb{N}$ . If  $\tau(p) = \tau(q)$  for any tracial state  $\tau$  of  $A$ , then there exist two contractions  $v_i$  in  $A \otimes M_n$ ,  $i = 0, 1$  such that*

$$\left\| \sum_{i=0,1} v_i^* v_i - p \otimes 1_n \right\| \leq \frac{4}{n}, \quad \left\| \sum_{i=0,1} v_i v_i^* - q \otimes 1_n \right\| \leq \frac{4}{n},$$

and

$$\text{dist}(v_i^* v_i, \{p \otimes x \mid x \in M_n\}) \leq \frac{2}{n} \quad \text{for } i = 0, 1.$$

*Proof.* Let  $e_j$ ,  $j = 1, 2, \dots, n$  be mutually orthogonal minimal projections of  $M_n$ . When  $n = 1, 2$ , or  $3$  the statement is trivial, thus we may assume that  $n \geq 4$ . Since  $A$  has strict comparison for projections, there exists a partial isometry  $r_1 \in A \otimes M_n$  such that

$$r_1^* r_1 = p \otimes e_1, \quad r_1 r_1^* \leq q \otimes (e_1 + e_2).$$

By the same reason, we obtain a partial isometry  $r_2 \in A \otimes M_n$  such that

$$r_2^* r_2 \leq p \otimes (e_2 + e_3), \quad r_2 r_2^* = q \otimes (e_1 + e_2) - r_1 r_1^*.$$

In the same way, there exists a partial isometry  $r_3 \in A \otimes M_n$  such that

$$r_3^* r_3 = p \otimes (e_2 + e_3) - r_2^* r_2, \quad r_3 r_3^* \leq q \otimes (e_3 + e_4).$$

Repeating this argument we have partial isometries  $r_j \in A \otimes M_n$ ,  $j = 1, 2, \dots, n-1$ . Set

$$f = \sum_{j=1}^n \frac{n-j}{n} e_j, \quad v_0 = \sum_{j=1}^{n-1} \sqrt{\frac{n-j}{n}} r_j.$$

Since  $\{r_j r_j^* \mid j = 1, 2, \dots, n-1\}$  and  $\{r_j^* r_j \mid j = 1, 2, \dots, n-1\}$  are sets of mutually orthogonal projections, it follows that

$$\|v_0^* v_0 - p \otimes f\| = \left\| \sum_{j=1}^{n-1} \frac{n-j}{n} r_j^* r_j - p \otimes f \right\| \leq \frac{2}{n}$$

and

$$\|v_0 v_0^* - q \otimes f\| = \left\| \sum_{j=1}^{n-1} \frac{n-j}{n} r_j r_j^* - q \otimes f \right\| \leq \frac{2}{n}.$$

Similarly we can construct a contraction  $v_1 \in A \otimes M_n$  such that

$$\|v_1^* v_1 - p \otimes (1_n - f)\| \leq \frac{2}{n}$$

and

$$\|v_1 v_1^* - q \otimes (1_n - f)\| \leq \frac{2}{n},$$

thereby completing the proof.  $\square$

### 3 Relative Property (SI)

In our previous work [18, Proposition 4.8], we showed certain strict comparison of central sequence algebras by using a technical condition named property (SI). The aim of this section is to introduce a relative version of property (SI) and to show a variant of [18, Proposition 4.8].

Until the end of this section, we let  $A$  and  $B$  be unital separable simple nuclear infinite-dimensional  $C^*$ -algebras and suppose that  $B$  has strict comparison. First, we recall the definitions of central sequence algebras. Let  $\omega$  be a free ultrafilter on  $\mathbb{N}$ , and let

$$\|b\|_2 = \max_{\tau \in T(B)} \tau(b^*b)^{1/2} \quad \text{for } b \in B,$$

$$c_\omega(B) = \left\{ (b_n)_n \in \ell^\infty(\mathbb{N}, B) \mid \lim_{n \rightarrow \omega} \|b_n\| = 0 \right\}.$$

We denote by  $B^\omega$  the ultraproduct  $C^*$ -algebra  $\ell^\infty(\mathbb{N}, B)/c_\omega(B)$ .

Set

$$J = \left\{ (b_n)_n \in \ell^\infty(\mathbb{N}, B) \mid \lim_{n \rightarrow \omega} \|b_n\|_2 = 0 \right\}$$

and  $M = \ell^\infty(\mathbb{N}, B)/J$ . Since  $c_\omega(B) \subset J$  we regard  $M$  as a quotient algebra of  $B^\omega$ , and denote by  $\Phi$  the quotient map from  $B^\omega$  to  $M$ . Note that if  $B$  has finitely many extreme traces then  $M$  is isomorphic to a finite direct sum of  $\text{II}_1$ -factors.

Let  $\pi$  be a unital embedding of  $A$  into  $B^\omega$ . We consider the relative commutants

$$\mathcal{B} = B^\omega \cap \pi(A)' \quad \text{and} \quad \mathcal{M} = M \cap \Phi(\pi(A))'.$$

The Choi-Effros lifting theorem allows us to get a sequence of unital completely positive maps  $\pi_n : A \rightarrow B$  such that  $(\pi_n(a))_n = \pi(a)$  in  $B^\omega$  for any  $a \in A$ .

The following theorem is based on the technique of Kirchberg and Rørdam [10, Theorem 3.3]. Combining this result and a generalized property (SI) (Lemma 3.2), we shall show the strict comparison of  $\mathcal{B}$  for projections in Proposition 3.3.

**Theorem 3.1** (Kirchberg-Rørdam). *The restriction  $\Phi|_{\mathcal{B}} : \mathcal{B} \rightarrow \mathcal{M}$  is surjective.*

*Proof.* Let  $x$  be a contraction in  $\mathcal{M}$ , and  $(b_n)_n \in B^\omega$  be such that  $\Phi((b_n)_n) = x$ . Set

$$D = C^*(\pi(A) \cup \{(b_n)_n\}) \subset B^\omega \quad \text{and} \quad I = \ker(\Phi|_D) \subset D.$$

Since  $x \in \Phi(\pi(A))'$  it follows that  $[\pi(a), (b_n)_n] \in I$  for any  $a \in A$ . Let  $e_\lambda \in I$ ,  $\lambda \in \Lambda$  be a quasicontral approximate unit of  $I$ . Then we have

$$\begin{aligned} 0 &= \lim_{\lambda \rightarrow \infty} \|(1 - e_\lambda)[(b_n)_n, \pi(a)](1 - e_\lambda)\| \\ &= \lim_{\lambda \rightarrow \infty} \|[(1 - e_\lambda)(b_n)_n(1 - e_\lambda), \pi(a)]\| \quad \text{for any } a \in A. \end{aligned}$$

This implies that for any finite subset  $F$  of  $A$  and  $\varepsilon > 0$  there exists a sequence  $e_{F, \varepsilon, n}$ ,  $n \in \mathbb{N}$  of positive contractions in  $B$  such that

$$\lim_{n \rightarrow \omega} \|e_{F, \varepsilon, n}\|_2 = 0, \quad \lim_{n \rightarrow \omega} \|[(1 - e_{F, \varepsilon, n})b_n(1 - e_{F, \varepsilon, n}), \pi_n(a)]\| < \varepsilon,$$

for all  $a \in F$ .

Let  $F_n$ ,  $n \in \mathbb{N}$  be an increasing sequence of finite subsets of  $A$  whose union is dense in  $A$  and  $\varepsilon_n > 0$ ,  $n \in \mathbb{N}$  a decreasing sequence which converges to 0. For any  $n \in \mathbb{N}$  we inductively obtain a subset  $\Omega_n \in \omega$  such that  $\Omega_{n+1} \subset \Omega_n$ ,  $\bigcap_{n \in \mathbb{N}} \Omega_n = \emptyset$  and

$$\|e_{F_n, \varepsilon_n, m}\|_2 < \varepsilon_n, \quad \|(1 - e_{F_n, \varepsilon_n, m})b_m(1 - e_{F_n, \varepsilon_n, m}), \pi_m(a)\| < \varepsilon_n,$$

for any  $m \in \Omega_n$  and  $a \in F_n$ . We define  $e_m = 0$  for  $m \in \mathbb{N} \setminus \Omega_1$  and  $e_m = e_{F_n, \varepsilon_n, m}$  for  $m \in \Omega_n \setminus \Omega_{n+1}$ . Then it follows that

$$\lim_{n \rightarrow \omega} \|e_n\|_2 = 0, \quad \lim_{n \rightarrow \omega} \|(1 - e_n)b_n(1 - e_n), \pi_n(a)\| = 0,$$

for any  $a \in \bigcup_{n \in \mathbb{N}} F_n$ . This means that

$$\Phi(((1 - e_n)b_n(1 - e_n))_n) = x, \quad ((1 - e_n)b_n(1 - e_n))_n \in \mathcal{B}.$$

□

**Lemma 3.2.** *Suppose that  $T(B)$  has a compact extreme boundary with finite topological dimension. Then  $B$  has property (SI) relative to  $\pi(A)$  in the following sense: For any sequences  $(e_n)_n$  and  $(f_n)_n$  of positive contractions in  $B$  satisfying  $(e_n)_n, (f_n)_n \in \mathcal{B}$ ,*

$$(e_n)_n \in J, \quad \lim_{m \rightarrow \infty} \lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau(f_n^m) > 0,$$

*there exists a sequence  $(s_n)_n$  in  $B$  such that  $(s_n)_n \in \mathcal{B}$ ,*

$$\lim_{n \rightarrow \omega} \|s_n^* s_n - e_n\| = 0, \quad \lim_{n \rightarrow \omega} \|f_n s_n - s_n\| = 0.$$

*Proof.* One can prove this lemma in a similar fashion to the proof of [17, Theorem 1.1]. We sketch a proof for the reader's convenience. In the proof we write  $x \approx_\varepsilon y$  if  $\|x - y\| < \varepsilon$  for  $x, y$  in a  $C^*$ -algebra.

Let  $F$  be a finite subset of contractions in  $A$  and  $\varepsilon > 0$ . Since  $A$  is nuclear there are two completely positive contractions  $\rho : A \rightarrow M_N$  and  $\sigma : M_N \rightarrow A$  such that  $\sigma \circ \rho(x) \approx_{\varepsilon/2} x$  for  $x \in F$ . Since  $A$  is unital simple infinite-dimensional, applying Voiculescu's theorem to  $\rho$  we can obtain  $d_i \in A$ ,  $i = 1, 2, \dots, N$  such that  $\rho(x) \approx_{\varepsilon/2} [\varphi(d_i^* x d_j)]_{i,j}$  for some pure state  $\varphi$  of  $A$  and for all  $x \in F$ . Identifying  $\sigma$  with a positive element of  $M_N(A)$  we have

$c_{l,i} \in A$ ,  $l, i = 1, 2, \dots, N$  such that  $\sigma(e_{i,j}) = \sum_{l=1}^N c_{l,i}^* c_{l,j}$  for the standard matrix units  $\{e_{i,j}\}_{i,j}$  of  $M_N$ . Thus it follows that

$$\sum_{l=1}^N \sum_{i,j=1}^N \varphi(d_i^* x d_j) \pi(c_{l,i}^* c_{l,j}) \approx_\varepsilon \pi(x) \quad \text{for } x \in F.$$

Due to the Akemann-Anderson-Pedersen excision theorem, there exists a positive contraction  $a \in A$  such that  $ad_i^* x d_j a \approx_{\varepsilon/N^2} \varphi(d_i^* x d_j) a^2$  for  $x \in F$  and  $\varphi(a) = 1$ . Because  $T(B)$  has compact extreme boundary with finite topological dimension, by [24, Theorem 1.1] (see also [10, 25]), we obtain a unital embedding of  $M_N$  into  $M \cap B'$ . By Theorem

3.1 we obtain a sequence of completely positive order zero maps  $\psi_n : M_N \rightarrow B$  such that  $\min_{\tau \in T(B)} \tau(\psi_n(1)^m) \rightarrow 1$  for any  $m \in \mathbb{N}$  and  $\|[\psi_n(x), b]\| \rightarrow 0$  as  $n \rightarrow \omega$  for any  $x \in M_N$  and  $b \in B$ . Since  $A$  is separable, we may further assume

$$\lim_{n \rightarrow \omega} \|[\psi_n(x), \pi_n(y)]\| = 0, \quad \lim_{n \rightarrow \omega} \|[\psi_n(x), f_n]\| = 0$$

for any  $x \in M_N$  and  $y \in A$ . Then a similar argument as in the proof of [17, Lemma 3.4] implies that there exist sequences  $(f_{l,n})_n$ ,  $l = 1, 2, \dots, N$  of positive contractions in  $B$  such that  $(f_{l,n})_n \in \mathcal{B}$ ,  $(f_{l,n}f_n)_n = (f_{l,n})_n$ ,  $(f_{l,n}f_{l',n})_n = 0$  for  $l \neq l'$  in  $B^\omega$ , and

$$\lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau(f_{l,n}^m) = \frac{1}{N} \lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau(f_n^m) \quad \text{for any } m \in \mathbb{N}.$$

Since  $A$  is unital and simple, there exist  $v_i \in A$ ,  $i = 1, 2, \dots, k$  such that  $\sum_{i=1}^k v_i^* a^2 v_i = 1$ .

Set  $\alpha = \left(\sum_{i=1}^k \|v_i\|^2\right)^{-1} > 0$ . By  $(f_{l,n})_n \in \mathcal{B}$  we see that for any  $m \in \mathbb{N}$

$$\lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau(f_{l,n}^{m/2} \pi_n(a)^2 f_{l,n}^{m/2}) \geq \alpha \lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau(f_{l,n}^m) > 0.$$

As  $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau(f_{l,n}^m) > 0$ ,  $\|a\| = 1$  and  $B$  has strict comparison, there exist sequences  $r_{l,n} \in B$ ,  $l = 1, 2, \dots, N$ ,  $n \in \mathbb{N}$  such that

$$\lim_{n \rightarrow \omega} \left\| r_{l,n}^* f_{l,n}^{1/2} \pi_n(a)^2 f_{l,n}^{1/2} r_{l,n} - e_n \right\| = 0, \quad \lim_{n \rightarrow \omega} \|r_{l,n}\| = \lim_{n \rightarrow \omega} \|e_n\|^{1/2} \leq 1.$$

Define

$$s_n = \sum_{l=1}^N \sum_{i=1}^N \pi_n(d_i a) f_{l,n}^{1/2} r_{l,n} \pi_n(c_{l,i}) \in B, \quad \text{for } n \in \mathbb{N}.$$

Then we have  $(f_n s_n)_n = (s_n)_n$  in  $B^\omega$ . And there exists a neighborhood  $\Omega \in \omega$  such that for  $x \in F$  and  $n \in \Omega$

$$\begin{aligned} s_n^* \pi_n(x) s_n &\approx \sum_{l=1}^N \sum_{i,j=1}^N \pi_n(c_{l,i}^*) r_{l,n}^* f_{l,n}^{1/2} \pi_n(a d_i^* x d_j a) f_{l,n}^{1/2} r_{l,n} \pi_n(c_{l,j}) \\ &\approx_\varepsilon \sum_l \sum_{i,j} \varphi(d_i^* x d_j) \pi_n(c_{l,i}^*) r_{l,n}^* f_{l,n}^{1/2} \pi_n(a^2) f_{l,n}^{1/2} r_{l,n} \pi_n(c_{l,j}) \\ &\approx \sum_l \sum_{i,j} \varphi(d_i^* x d_j) \pi_n(c_{l,i}^* c_{l,j}) e_n \approx_\varepsilon \pi_n(x) e_n. \end{aligned}$$

Since  $F \subset A$  and  $\varepsilon > 0$  are arbitrary,  $A$  is separable, and  $\pi$  is unital, we can find another sequence  $s_n$ ,  $n \in \mathbb{N}$  of  $B$  such that  $(f_n s_n)_n = (s_n)_n$ ,  $(s_n^* s_n)_n = (e_n)_n$  in  $B^\omega$  and  $(s_n)_n \in \mathcal{B}$ .  $\square$

**Proposition 3.3.** *Suppose that  $B$  has finitely many extremal tracial states. Then the following holds.*

- (i) *The map  $T(\mathcal{M}) \rightarrow T(\mathcal{B})$  induced by  $\Phi : \mathcal{B} \rightarrow \mathcal{M}$  is surjective.*

(ii)  $\mathcal{B}$  has strict comparison for projections.

*Proof of (i).* It suffices to show  $\tau(x) = 0$  for any  $\tau \in T(\mathcal{B})$  and  $x \in \mathcal{B} \cap \ker \Phi$ . We may assume that  $x$  is a positive contraction. Let  $(x_n)_n$  be a representative sequence of  $x$  consisting of positive contractions. We have

$$\lim_{n \rightarrow \omega} \min_{\tau \in T(B)} \tau((1 - x_n)^m) = 1$$

for any  $m \in \mathbb{N}$ . By Lemma 3.2,  $B$  has property (SI) relative to  $\pi(A)$ . It follows that there exists  $s_1 \in \mathcal{B} \cap \ker \Phi$  such that  $s_1^* s_1 = x$  and  $(1 - x)s_1 = s_1$ . Clearly  $x + s_1 s_1^*$  is a positive contraction in  $\mathcal{B} \cap \ker \Phi$ . Hence, in the same way we obtain  $s_2 \in \mathcal{B} \cap \ker \Phi$  such that  $s_2^* s_2 = x$  and  $(1 - x - s_1 s_1^*)s_2 = s_2$ . Then  $x + s_1 s_1^* + s_2 s_2^*$  is again a positive contraction in  $\mathcal{B} \cap \ker \Phi$ . Repeating this argument, one can find  $s_i \in \mathcal{B} \cap \ker \Phi$ ,  $i \in \mathbb{N}$  satisfying

$$s_i^* s_i = x \quad \text{and} \quad x + s_1 s_1^* + s_2 s_2^* + \cdots + s_m s_m^* \leq 1$$

for every  $m \in \mathbb{N}$ . Hence  $\tau(x) = 0$  for all  $\tau \in T(\mathcal{B})$ .  $\square$

*Proof of (ii).* Let  $\pi^{(n)}$  be the amplification map of  $\pi$  defined by  $\pi^{(n)}(a) = \pi(a) \otimes 1_n \in B^\omega \otimes M_n \cong (B \otimes M_n)^\omega$ ,  $n \in \mathbb{N}$ ,  $a \in A$ . By the canonical identification we have  $\mathcal{B} \otimes M_n \cong (B \otimes M_n)^\omega \cap \pi^{(n)}(A)'$  for any  $n \in \mathbb{N}$ . Then it suffices to consider two projections in  $\mathcal{B}$ .

Assume that two projections  $p$  and  $q$  in  $\mathcal{B}$  satisfy  $\tau(p) < \tau(q)$  for any  $\tau \in T(\mathcal{B})$ . This implies  $\bar{\tau} \circ \Phi(p) < \bar{\tau} \circ \Phi(q)$  for any  $\bar{\tau} \in T(\mathcal{M})$ . Because  $\mathcal{M}$  is a finite von Neumann algebra, there exists  $\bar{r} \in \mathcal{M}$  such that  $\bar{r}^* \bar{r} = \Phi(p)$  and  $\bar{r} \bar{r}^* \leq \Phi(q)$ . By Theorem 3.1 we obtain  $r \in \mathcal{B}$  such that  $\Phi(r) = \bar{r}$  and  $\|r\| \leq 1$ . Set  $w = qrp \in \mathcal{B}$  and let  $q_n, w_n \in B$ ,  $n \in \mathbb{N}$  are representatives of  $q$  and  $w$ . Since  $q$  is a projection and  $q \geq ww^*$ , we may assume that  $q_n, w_n \in B$  are projections and  $q_n \geq w_n w_n^*$  for any  $n \in \mathbb{N}$ . Since  $\tau(q - ww^*) \geq \tau(q - p) > 0$  for any  $\tau \in T(\mathcal{B})$  it follows that

$$\lim_{n \rightarrow \omega} \tau'(q_n - w_n w_n^*) > 0 \quad \text{for any } \tau' \in T(B).$$

Because  $\Phi(q - ww^*)^m = \Phi(q - ww^*)$  for any  $m \in \mathbb{N}$  and  $B$  has finitely many extremal traces, we have

$$\lim_{n \rightarrow \omega} \min_{\tau' \in T(B)} \tau'((q_n - w_n w_n^*)^m) = \lim_{n \rightarrow \omega} \min_{\tau' \in T(B)} \tau'(q_n - w_n w_n^*) > 0,$$

for any  $m \in \mathbb{N}$ . By Lemma 3.2 and  $p - w^* w \in \mathcal{B} \cap \ker \Phi$ , there exists  $s \in \mathcal{B} \cap \ker \Phi$  such that  $s^* s = p - w^* w$  and  $(q - ww^*)s = s$ . We define  $v = w + s$ , then we conclude that  $v^* v = w^* w + s^* s = p$  and  $qv = v$ . This completes the proof.  $\square$

## 4 Decomposition rank

We denote by  $\mathcal{U}$  the universal UHF-algebra. In this section, we show that certain  $\mathcal{U}$ -absorbing  $C^*$ -algebras have finite decomposition rank. First let us recall the definition of decomposition rank.

**Definition 4.1** (Kirchberg-Winter [11, 27]). A separable  $C^*$ -algebra  $A$  is said to have *decomposition rank at most  $d \in \mathbb{N} \cup \{0\}$*  if there exist completely positive contractions  $\varphi_n : A \rightarrow \bigoplus_{i=0}^d F_{i,n}$ ,  $n \in \mathbb{N}$  and  $\psi_{i,n} : F_{i,n} \rightarrow A$ ,  $i = 0, 1, \dots, d$ ,  $n \in \mathbb{N}$ , where  $F_{i,n}$  are finite-dimensional  $C^*$ -algebras, such that  $\psi_{i,n}$  are order zero (disjointness preserving),  $\sum_{i=0}^d \psi_{i,n} : \bigoplus_{i=0}^d F_{i,n} \rightarrow A$  is contractive, and

$$\lim_{n \rightarrow \infty} \left\| \left( \sum_{i=0}^d \psi_{i,n} \right) \circ \varphi_n(a) - a \right\| = 0 \quad \text{for any } a \in A,$$

where we simply write  $\sum_{i=0}^d \psi_{i,n}(\bigoplus_{i=0}^d x_i) = \sum_{i=0}^d \psi_{i,n}(x_i)$  for  $x_i \in F_{i,n}$ .

The proof of the following theorem is based on the crucial technique which was developed by A. Connes and U. Haagerup in the proof of [6, Theorem 3.1, (d) $\Rightarrow$ (a)] and [7, Theorem 4.2].

**Theorem 4.2.** *Let  $A$  be a unital separable simple nuclear  $C^*$ -algebra with a unique tracial state. If  $A$  is quasidiagonal then the decomposition rank of  $A \otimes \mathcal{U}$  is at most one.*

**Lemma 4.3.** *Let  $A$  be a unital simple stably finite  $C^*$ -algebra which absorbs a UHF-algebra tensorially,  $A_1$  a  $C^*$ -subalgebra of  $A$ , and  $\omega$  a free ultrafilter on  $\mathbb{N}$ . Suppose that  $v_n$ ,  $n \in \mathbb{N}$  are contractions in  $A$  such that*

$$(|v_n|)_n \in A_1^\omega \subset A^\omega.$$

*Then there exists a sequence  $(\tilde{v}_n)_n$  of contractions in  $A$  such that  $|\tilde{v}_n| \in A_1$  for any  $n \in \mathbb{N}$  and*

$$(\tilde{v}_n)_n = (v_n)_n \quad \text{in } A^\omega.$$

*Proof.* Let  $\varepsilon_n > 0$ ,  $n \in \mathbb{N}$  be a decreasing sequence which converges to 0, and let  $f_n(t) = \max\{0, t - \varepsilon_n\}$  for  $t \in [0, \infty)$ ,  $n \in \mathbb{N}$ . M. Rørdam showed that any unital simple stably finite UHF-absorbing  $C^*$ -algebra has stable rank one [21, Corollary 6.6]. Then, by G. K. Pedersen's polar decomposition theorem [19, Theorem 5], there exists a sequence  $u_n$ ,  $n \in \mathbb{N}$  of unitaries in  $A$  such that

$$\|u_n f_n(|v_n|) - v_n\| \leq \varepsilon_n.$$

By  $(|v_n|)_n \in A_1^\omega$  there exists a sequence  $h_n$ ,  $n \in \mathbb{N}$  of positive contractions in  $A_1$  such that  $(h_n)_n = (|v_n|)_n = (f_n(|v_n|))_n$  in  $A_1^\omega$ . Define  $\tilde{v}_n = u_n h_n$ . Then  $|\tilde{v}_n| = h_n \in A_1$  and  $(\tilde{v}_n)_n = (u_n f_n(|v_n|))_n = (v_n)_n$  in  $A^\omega$ .  $\square$

In what follows, we let  $\text{Ad } a(b) = aba^*$  for  $a, b$  in a  $C^*$ -algebra and denote by  $\|a\|_{2,\tau}$  the 2-norm of  $a$  associated with a tracial state  $\tau$ , i.e.  $\|a\|_{2,\tau} = \tau(a^*a)^{1/2}$ .

*Proof of Theorem 4.2.* Let  $\iota$  be the canonical unital embedding of  $A$  into  $A \otimes \mathcal{U}$ . Since  $A \otimes M_n$  also satisfies the assumptions in the theorem for any  $n \in \mathbb{N}$ , it suffices to construct completely positive contractions  $\tilde{\rho}_n : A \rightarrow M_{k_n} \oplus M_{k_n}$ ,  $n \in \mathbb{N}$  and  $\tilde{\sigma}_{i,n} : M_{k_n} \rightarrow A \otimes \mathcal{U}$ ,  $i = 0, 1$ ,  $n \in \mathbb{N}$  such that  $\tilde{\sigma}_{i,n}$  are order zero,  $\sum_{i=0,1} \tilde{\sigma}_{i,n} : M_{k_n} \oplus M_{k_n} \rightarrow A \otimes \mathcal{U}$  is contractive, and

$$\left\| \left( \sum_{i=0,1} \tilde{\sigma}_{i,n} \right) \circ \tilde{\rho}_n(a) - \iota(a) \right\| \rightarrow 0, \quad n \rightarrow \infty, \quad \text{for any } a \in A.$$

We let  $\mathcal{U}_i$ ,  $i = 0, 1$  be commuting unital  $C^*$ -subalgebras of  $\mathcal{U}$  such that  $\mathcal{U}_0 \cong \mathcal{U}_1 \cong \mathcal{U}$  and  $C^*(\mathcal{U}_0 \cup \mathcal{U}_1) = \mathcal{U}$ . We denote by  $\tau_A$ ,  $\tau_{\mathcal{U}_i}$ ,  $i = 0, 1$ , and  $\tau_{\mathcal{U}}$  the unique tracial states of  $A$ ,  $\mathcal{U}_i$ , and  $\mathcal{U}$ . Let  $\omega$  be a free ultrafilter on  $\mathbb{N}$ .

Since  $A$  is quasidiagonal, there are unital completely positive maps  $\rho_n : A \rightarrow M_{k_n}$ ,  $n \in \mathbb{N}$  such that

$$\lim_{n \rightarrow \infty} \|\rho_n(a)\rho_n(b) - \rho_n(ab)\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|\rho_n(a)\| = \|a\|,$$

for any  $a, b \in A$  (see [3, Lemma 7.1.4] for example). Let  $\sigma_n : M_{k_n} \rightarrow \mathcal{U}_0$ ,  $n \in \mathbb{N}$  be a sequence of unital embeddings. We define unital completely positive maps  $\varphi_n : A \rightarrow \mathcal{U}_0$ ,  $n \in \mathbb{N}$  and  $\varphi : A \rightarrow \mathcal{U}_0^\omega$  by

$$\varphi_n(a) = \sigma_n \circ \rho_n(a) \quad \text{and} \quad \varphi(a) = (\varphi_n(a))_n,$$

for any  $n \in \mathbb{N}$  and  $a \in A$ . Note that  $\varphi$  is a unital embedding because  $\rho_n$  are asymptotically multiplicative.

In what follows, we identify  $A \otimes \mathcal{U}_0$  with the set of constant sequences in  $(A \otimes \mathcal{U}_0)^\omega$  and  $\mathcal{U}_0$  with the unital  $C^*$ -subalgebra  $1_A \otimes \mathcal{U}_0$  of  $A \otimes \mathcal{U}_0$ . By this canonical unital embedding we also identify  $\mathcal{U}_0^\omega$  with the unital  $C^*$ -subalgebra of  $(A \otimes \mathcal{U}_0)^\omega$ . From these identifications it follows that  $[\iota(a), \varphi(b)] = 0$  for  $a, b \in A$ . We define the product map  $\iota \times \varphi : A \odot A \rightarrow (A \otimes \mathcal{U}_0)^\omega$  by  $(\iota \times \varphi)(a \otimes b) = \iota(a) \cdot \varphi(b)$  for  $a, b \in A$ , where  $\odot$  means the algebraic tensor product. By the nuclearity of  $A$  we can extend  $\iota \times \varphi$  to  $A \otimes A$ . Since  $\tau_A$  is the unique tracial state of  $A$ ,  $A \otimes A$  also has the unique tracial state  $\tau_A \otimes \tau_A$ . Then the strong closure of  $A \otimes A$  in the GNS-representation, associated with  $\tau_A \otimes \tau_A$ , is the injective  $\text{II}_1$ -factor, and the flip automorphism on  $A \otimes A$  is approximately inner in the strong topology [6], i.e., there exists a sequence  $w_n$ ,  $n \in \mathbb{N}$  of unitaries in  $A \otimes A$  such that

$$\lim_{n \rightarrow \infty} \|w_n(a \otimes b)w_n^* - b \otimes a\|_{2, \tau_A \otimes \tau_A} = 0,$$

for any  $a, b \in A$ . Set  $U_n = (\iota \times \varphi)(w_n) \in (A \otimes \mathcal{U}_0)^\omega$ ,  $n \in \mathbb{N}$ . We define the tracial state on  $(A \otimes \mathcal{U}_0)^\omega$  by  $(\tau_A \otimes \tau_{\mathcal{U}_0})_\omega((x_n)_n) = \lim_{n \rightarrow \omega} (\tau_A \otimes \tau_{\mathcal{U}_0})(x_n)$  for  $(x_n)_n \in (A \otimes \mathcal{U}_0)^\omega$ . Since  $\tau_A \otimes \tau_A$  is a unique tracial state of  $A \otimes A$  it follows that  $(\tau_A \otimes \tau_{\mathcal{U}_0})_\omega \circ (\iota \times \varphi)(x) = (\tau_A \otimes \tau_A)(x)$  for  $x \in A \otimes A$ , which implies that

$$\lim_{n \rightarrow \infty} \|U_n \varphi(a) U_n^* - \iota(a)\|_{2, (\tau_A \otimes \tau_{\mathcal{U}_0})_\omega} = 0,$$

for any  $a \in A$ . Since  $A$  is separable, by the standard reindexation trick we obtain a sequence  $u_n$ ,  $n \in \mathbb{N}$  of unitaries in  $A \otimes \mathcal{U}_0$  such that

$$\lim_{n \rightarrow \omega} \|u_n \varphi(a) u_n^* - \iota(a)\|_{2, \tau_A \otimes \tau_{\mathcal{U}_0}} = 0 \quad \text{for any } a \in A.$$

Set

$$\begin{aligned} \mathcal{A} &= (A \otimes \mathcal{U} \otimes M_2)^\omega, \\ \mathcal{A}_0 &= (A \otimes \mathcal{U}_0 \otimes M_2)^\omega \subset \mathcal{A}, \\ J_0 &= \{(x_n)_n \in \ell^\infty(A \otimes \mathcal{U}_0 \otimes M_2) \mid \lim_{n \rightarrow \omega} \|x_n\|_{2, \tau_A \otimes \tau_{\mathcal{U}_0} \otimes \text{tr}_2} = 0\}, \\ M_0 &= \ell^\infty(A \otimes \mathcal{U}_0 \otimes M_2) / J_0, \end{aligned}$$

and let  $\Phi_0$  be the quotient map from  $\mathcal{A}_0$  to  $M_0$ . For any  $a \in A$ , we define

$$\pi(a) = \left( \begin{bmatrix} \varphi_n(a) & 0 \\ 0 & \iota(a) \end{bmatrix} \right)_n \in \mathcal{A}_0, \quad u = \left( \begin{bmatrix} 0 & 0 \\ u_n & 0 \end{bmatrix} \right)_n \in \mathcal{A}_0.$$

Note that  $\pi$  is a unital embedding of  $A$  into  $\mathcal{A}_0$ . Because of  $\lim_{n \rightarrow \infty} \|u_n \varphi_n(a) - \iota(a) u_n\|_{2, \tau_A \otimes \tau_{\mathcal{U}_0}} = 0$  for any  $a \in A$ , we have

$$\Phi_0(u) \in M_0 \cap \Phi_0(\pi(A))'.$$

We also define

$$p = \begin{bmatrix} 1_{A \otimes \mathcal{U}_0} & 0 \\ 0 & 0 \end{bmatrix} \in \mathcal{A}_0, \quad q = \begin{bmatrix} 0 & 0 \\ 0 & 1_{A \otimes \mathcal{U}_0} \end{bmatrix} \in \mathcal{A}_0.$$

Then  $p$  and  $q$  belong to  $\mathcal{A}_0 \cap \pi(A)'$ . Since  $u_n$  are unitaries it follows that

$$u^* u = p \quad \text{and} \quad u u^* = q \quad \text{in } \mathcal{A}_0.$$

As  $\Phi_0(u)$ ,  $\Phi_0(p)$ , and  $\Phi_0(q)$  are in  $M_0 \cap \Phi_0(\pi(A))'$  we have

$$\bar{\tau} \circ \Phi_0(p) = \bar{\tau} \circ \Phi_0(q),$$

for any tracial state  $\bar{\tau}$  of  $M_0 \cap \Phi_0(\pi(A))'$ . Applying Proposition 3.3 (i) to  $B = A \otimes \mathcal{U}_0 \otimes M_2$ , we have

$$\tau(p) = \tau(q),$$

for any tracial state  $\tau$  of  $\mathcal{A}_0 \cap \pi(A)'$ .

By (ii) of Proposition 3.3,  $\mathcal{A}_0 \cap \pi(A)'$  has strict comparison for projections. Then by Lemma 2.1, for each  $n \in \mathbb{N}$  we obtain two contractions  $\tilde{w}_{i,n}$ ,  $i = 0, 1$  in  $(\mathcal{A}_0 \cap \pi(A))' \otimes M_n$  such that

$$\left\| \sum_{i=0,1} \tilde{w}_{i,n}^* \tilde{w}_{i,n} - p \otimes 1_n \right\| \leq \frac{4}{n}, \quad \left\| \sum_{i=0,1} \tilde{w}_{i,n} \tilde{w}_{i,n}^* - q \otimes 1_n \right\| \leq \frac{4}{n},$$

$$\text{and} \quad \text{dist}(\tilde{w}_{i,n}^* \tilde{w}_{i,n}, \{p \otimes x \mid x \in M_n\}) \leq \frac{2}{n} \quad \text{for } i = 0, 1.$$

We may think of  $M_n$  as a unital subalgebra of  $\mathcal{U}_1$ , and so for each  $n \in \mathbb{N}$  we get

$$\left\| \sum_{i=0,1} \tilde{w}_{i,n}^* \tilde{w}_{i,n} - p \right\| \leq \frac{4}{n}, \quad \left\| \sum_{i=0,1} \tilde{w}_{i,n} \tilde{w}_{i,n}^* - q \right\| \leq \frac{4}{n},$$

$$\text{and} \quad \text{dist}(\tilde{w}_{i,n}^* \tilde{w}_{i,n}, p(1_A \otimes \mathcal{U}_1 \otimes 1_2)^\omega) \leq \frac{2}{n} \quad \text{for } i = 0, 1.$$

Therefore one can find two sequences  $(\tilde{v}_{i,n})_n$ ,  $i = 0, 1$  of contractions in  $A \otimes \mathcal{U} \otimes M_2$  such that  $(\tilde{v}_{i,n})_n \in \mathcal{A} \cap \pi(A)'$  for  $i = 0, 1$ ,

$$\left( \sum_{i=0,1} \tilde{v}_{i,n}^* \tilde{v}_{i,n} \right)_n = p, \quad \left( \sum_{i=0,1} \tilde{v}_{i,n} \tilde{v}_{i,n}^* \right)_n = q \quad \text{in } \mathcal{A},$$

and  $(|\tilde{v}_{i,n}|)_n \in p(1_A \otimes \mathcal{U}_1 \otimes 1_2)^\omega \subset p\mathcal{A}p$ , for  $i = 0, 1$ .

Hence there exist two sequences  $(v_{i,n})_n$ ,  $i = 0, 1$  of contractions in  $A \otimes \mathcal{U}$  such that

$$\left( \begin{bmatrix} 0 & 0 \\ v_{i,n} & 0 \end{bmatrix} \right)_n = (\tilde{v}_{i,n})_n \in \mathcal{A}, \quad i = 0, 1,$$

and

$$\left( \sum_{i=0,1} v_{i,n}^* v_{i,n} \right)_n = \left( \sum_{i=0,1} v_{i,n} v_{i,n}^* \right)_n = 1_{A \otimes \mathcal{U}} \quad \text{in } (A \otimes \mathcal{U})^\omega.$$

Because of  $(\tilde{v}_{i,n})_n \in \mathcal{A} \cap \pi(A)'$  we have

$$(v_{i,n} \varphi_n(a))_n = (\iota(a) v_{i,n})_n \quad \text{in } (A \otimes \mathcal{U})^\omega,$$

for any  $a \in A$ . It follows from these conditions that for any  $a \in A$

$$\begin{aligned} \iota(a) &= \sum_{i=0,1} (v_{i,n} \varphi_n(a) v_{i,n}^*)_n \\ &= \sum_{i=0,1} (\text{Ad } v_{i,n} \circ \sigma_n \circ \rho_n(a))_n \quad \text{in } (A \otimes \mathcal{U})^\omega. \end{aligned}$$

From  $(|\tilde{v}_{i,n}|)_n \in p(1_A \otimes \mathcal{U}_1 \otimes 1_2)^\omega$  we have  $(|v_{i,n}|)_n \in (1_A \otimes \mathcal{U}_1)^\omega \subset (A \otimes \mathcal{U})^\omega$ . By Lemma 4.3 we may assume that  $|v_{i,n}|$  is in  $1_A \otimes \mathcal{U}_1$  for every  $n \in \mathbb{N}$ . For  $i = 0, 1$  and  $n \in \mathbb{N}$ , we define an order zero completely positive contraction  $\tilde{\sigma}_{i,n} : M_{k_n} \rightarrow A \otimes \mathcal{U}$ ,  $n \in \mathbb{N}$  by  $\tilde{\sigma}_{i,n} = \text{Ad } v_{i,n} \circ \sigma_n$  and simply write  $\sum_{i=0,1} \tilde{\sigma}_{i,n}(x_0 \oplus x_1) = \sum_{i=0,1} \tilde{\sigma}(x_i)$  for  $x_i \in M_{k_n}$ . Since  $\iota$  and  $\rho_n$  are unital it follows that

$$\lim_{n \rightarrow \omega} \left\| \sum_{i=0,1} \tilde{\sigma}_{i,n} \right\| = 1.$$

By multiplying  $\tilde{\sigma}_{i,n}$  by suitable scalars, we may further assume that  $\sum_{i=0,1} \tilde{\sigma}_{i,n}$  are contractive. Define completely positive contractions  $\tilde{\rho}_n$ ,  $n \in \mathbb{N}$  from  $A$  to  $M_{k_n} \oplus M_{k_n}$  by  $\tilde{\rho}_n(a) = \rho_n(a) \oplus \rho_n(a)$ . Then we conclude that

$$\iota(a) = \left( \left( \sum_{i=0,1} \tilde{\sigma}_{i,n} \right) \circ \tilde{\rho}_n(a) \right)_n \quad \text{in } (A \otimes \mathcal{U})^\omega,$$

for any  $a \in A$ . This completes the proof.  $\square$

## 5 $\mathcal{Z}$ -absorbing $C^*$ -algebras

In this section, we prove Theorem 1.1.

*Proof of Theorem 1.1.* By Theorem [17, Theorem 1.1],  $A$  is  $\mathcal{Z}$ -absorbing, i.e.  $A \cong A \otimes \mathcal{Z}$ . Let  $F$  be a finite subset of contractions in  $A$  and let  $\varepsilon > 0$ . By Theorem 4.2, decomposition rank of  $A \otimes \mathcal{U}$  is at most one, and so there exist completely positive contractions  $\varphi : A \rightarrow$

$E_0 \oplus E_1$  and  $\psi_i : E_i \rightarrow A \otimes \mathcal{U}$ ,  $i = 0, 1$  such that  $E_i$  are finite dimensional  $C^*$ -algebras,  $\psi_i$  are order zero,  $\sum_{i=0,1} \psi_i : E_0 \oplus E_1 \rightarrow A \otimes \mathcal{U}$  is contractive and

$$\left\| \left( \sum_{i=0,1} \psi_i \right) \circ \varphi(a) - a \otimes 1_{\mathcal{U}} \right\| < \varepsilon/2$$

for any  $a \in F$ . Any order zero completely positive map from a finite dimensional  $C^*$ -algebra can be identified with a  $*$ -homomorphism from the cone over the algebra (see [27, Proposition 1.2] for instance). A cone over a finite dimensional  $C^*$ -algebra is projective [15]. Therefore we may assume that  $\psi_i$ ,  $i = 0, 1$  satisfy  $\psi_i(E_i) \subset A \otimes M_l$  for some unital matrix subalgebra  $M_l$  of  $\mathcal{U}$ .

Choose  $m \in \mathbb{N}$  so that  $(m-1)^{-1} < \varepsilon/2$ . Let  $e_1, e_2, \dots, e_{lm} \in M_{lm}$  and  $f_0, f_1, \dots, f_{lm} \in M_{lm+1}$  be mutually orthogonal minimal projections. For  $j = 1, 2, \dots, m$ , we put

$$I_j = \{n \in \mathbb{N} \mid (j-1)l + 1 \leq n \leq jl\}.$$

Let  $\alpha_j : M_l \rightarrow M_{lm}$ ,  $\beta_j : M_l \rightarrow M_{lm+1}$  and  $\gamma_j : M_l \rightarrow M_{lm+1}$  be embeddings such that

$$\alpha_j(1_l) = \sum_{n \in I_j} e_n, \quad \beta_j(1_l) = \sum_{n \in I_j} f_n, \quad \gamma_j(1_l) = \sum_{n \in I_j} f_{n-1}.$$

Take a unitary  $w \in M_l \otimes M_l$  such that  $w(x \otimes y)w^* = y \otimes x$  for any  $x, y \in M_l$ . Let  $u : [0, 1] \rightarrow M_l \otimes M_l$  be a continuous path of unitaries such that  $u(0) = 1_l$  and  $u(1) = w$ . Define a continuous path of unitaries  $\tilde{u} : [0, 1] \rightarrow M_{lm} \otimes M_{lm+1}$  by

$$\tilde{u}(t) = 1_{lm} \otimes f_0 + \sum_{j,k=1}^m (\alpha_j \otimes \beta_k)(u(t)).$$

For  $i = 0, 1$ , we define order zero completely positive maps  $\lambda_i : M_l \rightarrow C([0, 1]) \otimes M_{lm} \otimes M_{lm+1}$  by

$$\lambda_0(x)(t) = \tilde{u}(t) \left( \sum_{j=1}^m (1-t) \alpha_j(x) \otimes f_0 + \sum_{j,k=1}^m \left( 1 - \frac{(m-k)t}{m-1} \right) \alpha_j(x) \otimes \beta_k(1_l) \right) \tilde{u}(t)^*$$

and

$$\lambda_1(x)(t) = \sum_{k=1}^m \frac{(m-k)t}{m-1} 1_{lm} \otimes \gamma_k(x).$$

It is straightforward to check

$$1 - \frac{t}{m-1} \leq \lambda_0(1_l)(t) + \lambda_1(1_l)(t) \leq 1$$

for any  $t \in [0, 1]$ . This means that  $\lambda_0 + \lambda_1$  is contractive. Moreover one has

$$\lambda_0(x)(0) = \sum_{j=1}^m \alpha_j(x) \otimes 1_{lm+1}, \quad \lambda_0(x)(1) = \sum_{k=1}^m \left( 1 - \frac{m-k}{m-1} \right) 1_{lm} \otimes \beta_k(x)$$

and

$$\lambda_1(x)(0) = 0, \quad \lambda_1(x)(1) = \sum_{k=1}^m \frac{m-k}{m-1} 1_{lm} \otimes \gamma_k(x).$$

Hence we may regard  $\lambda_i$  as a map from  $M_l$  to the dimension drop algebra

$$Z = \{g \in C([0, 1]) \otimes M_{lm} \otimes M_{lm+1} \mid g(0) \in M_{lm} \otimes \mathbb{C}, g(1) \in \mathbb{C} \otimes M_{lm+1}\}.$$

Then  $(\text{id}_A \otimes \lambda_i) \circ \psi_j$  is an order zero map from  $E_j$  to  $A \otimes Z$  for each  $i, j = 0, 1$ . We also obtain

$$\begin{aligned} & \left\| \left( \sum_{i,j} (\text{id}_A \otimes \lambda_i) \circ \psi_j \right) \circ (\varphi \oplus \varphi)(a) - a \otimes 1_Z \right\| \\ & < \left\| \sum_{i=0,1} \text{id}_A \otimes \lambda_i(a \otimes 1_l) - a \otimes 1_Z \right\| + \frac{\varepsilon}{2} \\ & \leq \frac{1}{m-1} + \frac{\varepsilon}{2} < \varepsilon \end{aligned}$$

for every  $a \in F$ . Since there exists a unital embedding  $Z \rightarrow \mathcal{Z}$ , we can conclude that  $A \cong A \otimes \mathcal{Z}$  has decomposition rank at most three.  $\square$

## 6 $C^*$ -algebras with tracial rank zero

W. Winter [26, 27] proved that if a unital separable simple  $C^*$ -algebra  $A$  has (locally) finite decomposition rank and has real rank zero, then  $A$  has tracial rank zero in the sense of H. Lin [12]. In this section, using the same strategy as in the proof of Theorem 4.2 we give a direct proof that  $A$  has tracial rank zero under the assumption that the  $C^*$ -algebra absorbs a UHF-algebra tensorially and has a unique tracial state.

**Theorem 6.1.** *Let  $A$  be a unital separable simple nuclear quasidiagonal  $C^*$ -algebra with a unique tracial state. Then the tracial rank of  $A \otimes B$  is zero for any UHF-algebra  $B$ .*

*Proof.* Because of [16, Lemma 2.4] it is enough to show that  $A \otimes \mathcal{U}$  has tracial rank zero. We use the same notations as in the proof of Theorem 4.2. Let  $\varepsilon > 0$ . Since  $A \otimes M_n$  also satisfies the same assumption in the theorem for any  $n \in \mathbb{N}$ , it suffices to construct a sequence of finite dimensional  $C^*$ -subalgebras  $B_n$ ,  $n \in \mathbb{N}$  of  $A \otimes \mathcal{U}$  such that

$$\tau_{A \otimes \mathcal{U}}(1_{A \otimes \mathcal{U}} - 1_{B_n}) < \varepsilon, \quad \lim_{n \rightarrow \omega} \|[1_{B_n}, \iota(a)]\| = 0,$$

and

$$\lim_{n \rightarrow \omega} \text{dist}(\iota(a)1_{B_n}, B_n) = 0, \quad \text{for any } a \in A.$$

Recall that  $\varphi_n : A \rightarrow \mathcal{U}_0$  is the composition of the unital completely positive map  $\rho_n : A \rightarrow M_{k_n}$  and the unital embedding  $\sigma_n : M_{k_n} \rightarrow \mathcal{U}_0$ . Clearly we have  $[\varphi_n(a), 1_A \otimes x] = 0$  for any  $n \in \mathbb{N}$ ,  $a \in A$ , and  $x \in \mathcal{U}_1$ . For any  $\tau \in T(\mathcal{A} \cap \pi(A)')$  we have seen that

$\tau(p) = \tau(q) = 1/2$  in the proof of Theorem 4.2 . Thus for any  $\tau \in T(\mathcal{A} \cap \pi(A)'),$  the functional

$$\mathcal{U}_1 \ni x \mapsto 2\tau \left( \left( \begin{bmatrix} 1_A \otimes x & 0 \\ 0 & 0 \end{bmatrix} \right)_n \right) \in \mathbb{C}$$

gives the unique tracial state  $\tau_{\mathcal{U}_1}$  of  $\mathcal{U}_1$ . Let  $e_0$  be a projection in  $\mathcal{U}_1$  such that  $1 - \varepsilon < \tau_{\mathcal{U}_1}(e_0) < 1$ . Set

$$e = \left( \begin{bmatrix} 1_A \otimes e_0 & 0 \\ 0 & 0 \end{bmatrix} \right)_n \in \mathcal{A} \cap \pi(A)'.$$

Then we have

$$\tau(e) = \frac{1}{2} \tau_{\mathcal{U}_1}(e_0) < \frac{1}{2} = \tau(p) = \tau(q) \quad \text{for any } \tau \in T(\mathcal{A} \cap \pi(A)').$$

Applying (ii) of Proposition 3.3 to  $B = A \otimes \mathcal{U} \otimes M_2$ , we see that  $\mathcal{A} \cap \pi(A)'$  has strict comparison for projections. We obtain a partial isometry  $\tilde{v} \in \mathcal{A} \cap \pi(A)'$  such that

$$\tilde{v}^* \tilde{v} = e \quad \text{and} \quad q \tilde{v} = \tilde{v}.$$

By the definition of  $p$  and  $q$  there exists a sequence  $v_n, n \in \mathbb{N}$  of contractions in  $A \otimes \mathcal{U}$  such that

$$\left( \begin{bmatrix} 0 & 0 \\ v_n & 0 \end{bmatrix} \right)_n = \tilde{v} \quad \text{in } \mathcal{A}, \quad \text{and} \quad (v_n^* v_n)_n = (1_A \otimes e_0)_n \quad \text{in } (A \otimes \mathcal{U})^\omega.$$

By perturbing  $v_n$  slightly, we may further assume  $v_n^* v_n = 1_A \otimes e_0$ . Because of  $\tilde{v} \in \pi(A)'$  we have

$$(v_n \varphi_n(a))_n = (\iota(a) v_n)_n \quad \text{in } (A \otimes \mathcal{U})^\omega \text{ for any } a \in A.$$

Then it follows that  $[(v_n v_n^*)_n, \iota(a)] = 0$  in  $(A \otimes \mathcal{U})^\omega$  for any  $a \in A$ . We define finite dimensional  $C^*$ -subalgebras  $B_n \subset A \otimes \mathcal{U}$  for  $n \in \mathbb{N}$  by

$$B_n = \text{Ad } v_n (\sigma_n(M_{k_n})).$$

One can verify that these  $B_n$  satisfy the desired properties. □

The following corollary is a direct consequence of [14, Theorem 5.4], [17, Theorem 1.1] and the theorem above.

**Corollary 6.2.** *Let  $A$  be a unital separable simple nuclear quasidiagonal  $C^*$ -algebra with a unique tracial state. Suppose that  $A$  has strict comparison and satisfies the UCT. Then  $A$  is classifiable by the Elliott invariants and is isomorphic to a unital simple approximately subhomogeneous algebra.*

## 7 Kirchberg algebras

A separable simple nuclear purely infinite  $C^*$ -algebra is called a Kirchberg algebra. In this section we prove the following theorem. See [28, Definition 2.1] for the definition of nuclear dimension.

**Theorem 7.1.** *Any Kirchberg algebra has nuclear dimension at most three.*

*Proof.* By [28, Corollary 2.8], it suffices to show that a unital Kirchberg algebra  $A$  has nuclear dimension at most three. Because of the Kirchberg-Phillips theorem,  $A$  is isomorphic to  $A \otimes \mathcal{O}_\infty$ . We let  $B_i$ ,  $i = 0, 1$  be commuting unital  $C^*$ -subalgebras of  $\mathcal{O}_\infty$  such that  $B_0 \cong B_1 \cong \mathcal{O}_\infty$  and  $C^*(B_0 \cup B_1) = \mathcal{O}_\infty$ . Let  $s \in B_0$  be a non-unitary isometry and put  $q = ss^*$ ,  $p = 1_{B_0} - q$ . Define  $\iota : A \rightarrow A \otimes B_0$  by  $\iota(a) = a \otimes q$ . The Kirchberg-Phillips theorem also tells us that there exists a unital embedding  $\rho : A \rightarrow \mathcal{O}_2$ . Since  $pB_0p$  contains a unital copy of  $\mathcal{O}_2$ , there exists a unital embedding  $\sigma : \mathcal{O}_2 \rightarrow 1_A \otimes pB_0p \subset A \otimes B_0$ . Let  $\varphi = \sigma \circ \rho$ .

We would like to apply the same argument as the proof of Theorem 4.2 to  $\varphi$  and  $\iota$ . Set

$$\mathcal{A} = (A \otimes \mathcal{O}_\infty)^\omega, \quad \mathcal{A}_0 = (A \otimes B_0)^\omega \subset \mathcal{A}.$$

Define a unital homomorphism  $\pi : A \rightarrow \mathcal{A}_0$  by

$$\pi(a) = \varphi(a) + \iota(a)$$

for  $a \in A$ . By [9, Corollary 4],  $\mathcal{A}_0 \cap \pi(A)'$  is purely infinite simple. In particular, for any non-zero projections  $e, f \in (\mathcal{A}_0 \cap \pi(A)') \otimes M_k$ ,  $e$  is Murray-von Neumann equivalent to a subprojection of  $f$ . Hence, for any  $n \in \mathbb{N}$ , we can apply (the proof of) Lemma 2.1 and obtain contractions

$$w_{i,n} \in (\mathcal{A}_0 \cap \pi(A)') \otimes (M_n \oplus M_{n+1}), \quad i = 0, 1,$$

such that

$$\left\| \sum_{i=0,1} w_{i,n}^* w_{i,n} - 1_A \otimes p \otimes (1_n \oplus 1_{n+1}) \right\| \leq \frac{4}{n}, \quad \left\| \sum_{i=0,1} w_{i,n} w_{i,n}^* - 1_A \otimes q \otimes (1_n \oplus 1_{n+1}) \right\| \leq \frac{4}{n},$$

and

$$\text{dist}(w_{i,n}^* w_{i,n}, \{1_A \otimes p \otimes x \mid x \in M_n \oplus M_{n+1}\}) \leq \frac{2}{n}.$$

Since  $M_n \oplus M_{n+1}$  can be unitaly embedded in  $B_1 \cong \mathcal{O}_\infty$  and  $n \in \mathbb{N}$  is arbitrary, there exist  $v_i \in \mathcal{A} \cap \pi(A)'$ ,  $i = 0, 1$  such that  $\sum_{i=0,1} v_i^* v_i = 1_A \otimes p$ ,  $\sum_{i=1,0} v_i v_i^* = 1_A \otimes q$ , and  $v_i^* v_i \in (1_A \otimes pB_1)^\omega$ ,  $i = 0, 1$ .

In the same way as the proof of Theorem 4.2, one has

$$\begin{aligned} a \otimes 1_{\mathcal{O}_\infty} &= (1_A \otimes s^*) \iota(a) (1_A \otimes s) \\ &= (1_A \otimes s^*) \left( \sum_{i=0,1} v_i \varphi(a) v_i^* \right) (1_A \otimes s) \\ &= \sum_{i=0,1} \text{Ad}(1_A \otimes s^*) \circ \text{Ad } v_i \circ \sigma \circ \rho(a) \quad \text{in } \mathcal{A}, \end{aligned}$$

for every  $a \in A$ . For  $i = 0, 1$ , we define an order zero completely positive contraction  $\tilde{\sigma}_i : \mathcal{O}_2 \rightarrow \mathcal{A}$  by  $\tilde{\sigma}_i = \text{Ad}(1_A \otimes s^*) \circ \text{Ad } v_i \circ \sigma$ .

Let  $F \subset A$  be a finite subset and let  $\varepsilon > 0$ . By [28, Theorem 7.4], the nuclear dimension of  $\mathcal{O}_2$  is one. It follows that there exist completely positive contractions  $\alpha : \mathcal{O}_2 \rightarrow E_0 \oplus E_1$  and  $\beta_j : E_j \rightarrow \mathcal{O}_2$ ,  $j = 0, 1$ , such that  $E_j$  are finite dimensional  $C^*$ -algebras,  $\beta_j$  are order zero and

$$\left\| \left( \sum_{j=0,1} \beta_j \right) \circ \alpha(\rho(a)) - \rho(a) \right\| < \varepsilon$$

for any  $a \in F$ . Then  $\tilde{\sigma}_i \circ \beta_j$  is an order zero map from  $E_j$  to  $\mathcal{A}$  for each  $i, j = 0, 1$ . Define a completely positive contraction  $\tilde{\rho} : A \rightarrow E_0 \oplus E_1 \oplus E_0 \oplus E_1$  by  $\tilde{\rho}(a) = \alpha(\rho(a)) \oplus \alpha(\rho(a))$ . We have

$$\left\| \left( \sum_{i,j=0,1} \tilde{\sigma}_i \circ \beta_j \right) \circ \tilde{\rho}(a) - a \otimes 1_{\mathcal{O}_\infty} \right\| < \varepsilon$$

for any  $a \in F$ . Since the cone over  $E_j$  is projective,  $\tilde{\sigma}_i \circ \beta_j$  lifts to an order zero map to  $\ell^\infty(\mathbb{N}, A \otimes \mathcal{O}_\infty)$ . The proof is completed.  $\square$

## References

- [1] C. A. Akemann, J. Anderson, and G. K. Pedersen, *Excising states of  $C^*$ -algebras*, Canad. J. Math. 38 (1986), 1239–1260.
- [2] B. Blackadar, *Comparison theory for simple  $C^*$ -algebras*, Operator algebras and applications, London Math. Soc. Lecture Note Ser., 135, Cambridge Univ. Press, Cambridge, (1988), 21–54.
- [3] N. P. Brown and N. Ozawa,  *$C^*$ -algebras and finite-dimensional approximations*, Graduate Studies in Mathematics, 88. Amer. Math. Soc., Providence, RI, 2008.
- [4] A. Connes, *Periodic automorphisms of the hyperfinite factor of type  $II_1$* , Acta Sci. Math. 39 (1977), no. 1-2, 39–66.
- [5] A. Connes, *Outer conjugacy classes of automorphisms of factors*, Ann. Sci. École Norm. Sup. (4) 8 (1975), no. 3, 383–419.
- [6] A. Connes, *Classification of injective factors: cases  $II_1$ ,  $II_\infty$ ,  $III_\lambda$ ,  $\lambda \neq 1$* , Ann. of Math. 104 (1976), 73–115.
- [7] U. Haagerup, *A new proof of the equivalence of injectivity and hyperfiniteness for factors on a separable Hilbert space*, J. Funct. Anal. 62 (1985), no. 2, 160–201.
- [8] X. Jiang and H. Su, *On a simple unital projectionless  $C^*$ -algebra*, Amer. J. Math. 121 (1999), 359–413.
- [9] E. Kirchberg, *The classification of purely infinite  $C^*$ -algebras using Kasparov’s theory*, preprint, 1994.
- [10] E. Kirchberg and M. Rørdam, *Central sequence  $C^*$ -algebras and tensorial absorption of the Jiang-Su algebra*, To appear in J. Reine Angew. Math.

- [11] E. Kirchberg and W. Winter, *Covering dimension and quasidiagonality*, Internat. J. Math. 15 (2004), no. 1, 63–85.
- [12] H. Lin, *An introduction to the classification of amenable  $C^*$ -algebras*, World Scientific Publishing Co., Inc., River Edge, NJ, 2001.
- [13] H. Lin, *Tracially AF  $C^*$ -algebras*, Trans. Amer. Math. Soc. 353 (2001), no. 2, 693–722.
- [14] H. Lin and Z. Niu, *Lifting  $KK$ -elements, asymptotical unitary equivalence and classification of simple  $C^*$ -algebras*, Adv. Math. 219 (2008), 1729–1769.
- [15] T. Loring, *Lifting solutions to perturbing problems in  $C^*$ -algebras*, Fields Institute monographs 8, AMS, Providence, Rhode Island (1997).
- [16] H. Matui and Y. Sato,  *$\mathcal{Z}$ -stability of crossed products by strongly outer actions*, Comm. Math. Phys. 314 (2012), no. 1, 193–228.
- [17] H. Matui and Y. Sato, *Strict comparison and  $\mathcal{Z}$ -absorption of nuclear  $C^*$ -algebras*, Acta Math. 209 (2012), no. 1, 179–196.
- [18] H. Matui and Y. Sato,  *$\mathcal{Z}$ -stability of crossed products by strongly outer actions II*, preprint, arXiv:1205.1590.
- [19] G. K. Pedersen, *Unitary extensions and polar decompositions in a  $C^*$ -algebra*, J. Operator Theory 17 (1987), no. 2, 357–364.
- [20] S. Popa, *A short proof of "injectivity implies hyperfiniteness" for finite von Neumann algebras*, J. Operator Theory 16 (1986), no. 2, 261–272.
- [21] M. Rørdam, *On the structure of simple  $C^*$ -algebras tensored with a UHF-algebra*, J. Funct. Anal. 100 (1991), no. 1, 1–17.
- [22] M. Rørdam, *The stable and the real rank of  $\mathcal{Z}$ -absorbing  $C^*$ -algebras*, Internat. J. Math. 15 (2004), no. 10, 1065–1084. arXiv:math/0408020.
- [23] M. Rørdam and W. Winter, *The Jiang-Su algebra revisited*, J. Reine. Angew. Math. 642 (2010), 129–155. arXiv:0801.2259.
- [24] Y. Sato, *Trace spaces of simple nuclear  $C^*$ -algebras with finite-dimensional extreme boundary*, preprint, arXiv:1209.3000.
- [25] A. S. Toms, S. White, W. Winter,  *$\mathcal{Z}$ -stability and finite dimensional tracial boundaries*, preprint, arXiv:1209.3292.
- [26] W. Winter, *Simple  $C^*$ -algebras with locally finite decomposition rank*, J. Funct. Anal. 243 (2007), no. 2, 394–425.
- [27] W. Winter, *Decomposition rank and  $\mathcal{Z}$ -stability* Invent. Math. 179 (2010), no. 2, 229–301.
- [28] W. Winter and J. Zacharias, *The nuclear dimension of  $C^*$ -algebras*, Adv. Math. 224 (2010), no. 2, 461–498, arXiv:0903.4914.